

CD4 Rabbit pAb

Catalog No.: A0362 **15 Publications**

Basic Information

Observed MW

51kDa

Calculated MW

51kDa

Category

Primary antibody

Applications

ELISA, WB, IF/ICC

Cross-Reactivity

Human, Mouse, Rat

Background

This gene encodes the CD4 membrane glycoprotein of T lymphocytes. The CD4 antigen acts as a coreceptor with the T-cell receptor on the T lymphocyte to recognize antigens displayed by an antigen presenting cell in the context of class II MHC molecules. The CD4 antigen is also a primary receptor for entry of the human immunodeficiency virus through interactions with the HIV Env gp120 subunit. This gene is expressed not only in T lymphocytes, but also in B cells, macrophages, granulocytes, as well as in various regions of the brain. The protein functions to initiate or augment the early phase of T-cell activation, and may function as an important mediator of indirect neuronal damage in infectious and immune-mediated diseases of the central nervous system. Multiple alternatively spliced transcript variants encoding different isoforms have been identified in this gene.

Recommended Dilutions

WB 1:500 - 1:2000**IF/ICC** 1:50 - 1:200

Immunogen Information

Gene ID

920

Swiss Prot

P01730

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 26-200 of human CD4 (NP_000607.1).

Synonyms

T4; IMD79; Leu-3; OKT4D; CD4mut; CD4

Contact

 | 400-999-6126 | cn.market@abclonal.com.cn | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal, 50% glycerol, pH7.3.


Validation Data




Western blot analysis of lysates from U-937 cells, using CD4 Rabbit pAb (A0362) at 1:500 dilution.
Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 30s.



Immunofluorescence analysis of Mouse thymus using CD4 Rabbit pAb (A0362) at dilution of 1:100. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of Rat thymus cells using CD4 Rabbit pAb (A0362) at dilution of 1:100. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of Jurkat cells using CD4 Rabbit pAb (A0362) at dilution of 1:100. Blue: DAPI for nuclear staining.