# **FAH Rabbit pAb**

Catalog No.: A13492 4 Publications



## **Basic Information**

### **Observed MW**

40-45kDa

### **Calculated MW**

46kDa

### Category

Primary antibody

### **Applications**

ELISA,WB

### **Cross-Reactivity**

Human, Mouse, Rat

## **Background**

Predicted to enable fumarylacetoacetase activity. Predicted to be involved in L-phenylalanine catabolic process; homogentisate catabolic process; and tyrosine catabolic process. Predicted to act upstream of or within arginine catabolic process. Located in extracellular exosome. Implicated in tyrosinemia type I.

## **Recommended Dilutions**

**WB** 

1:2000 - 1:8000

## **Immunogen Information**

Gene ID 2184 **Swiss Prot** 

P16930

### **Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 1-419 of human FAH (NP\_000128.1).

## **Synonyms**

FAH

## **Contact**

<b>a</b>		400-999-6126
$\bowtie$		cn.market@abclonal.com.cn
•	Ī	www.abclonal.com.cn


## **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

### **Storage**

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.



Western blot analysis of various lysates, using FAH Rabbit pAb (A13492) at 1:7000 dilution. Secondary antibody: HRP Goat Anti-Rabbit  $\lg G$  (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 60s.