

# OCIAD2 Rabbit pAb

Catalog No.: A14962

## Basic Information

### Observed MW

17kDa

### Calculated MW

17kDa

### Category

Primary antibody

### Applications

ELISA, WB, IF/ICC

### Cross-Reactivity

Human

## Background

Predicted to be involved in endocytosis; hematopoietic stem cell homeostasis; and positive regulation of receptor signaling pathway via JAK-STAT. Predicted to act upstream of or within response to bacterium. Predicted to be located in Golgi apparatus; endosome; and lysosome.

## Recommended Dilutions

WB 1:500 - 1:2000

IF/ICC 1:50 - 1:200

## Immunogen Information

### Gene ID

132299

### Swiss Prot

Q56VL3

### Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 1-99 of human OCIAD2 (NP\_689611.1).

### Synonyms

OCIAD2

## Contact

☎ | 400-999-6126

✉ | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn)

🌐 | [www.abclonal.com.cn](http://www.abclonal.com.cn)

## Product Information

### Source

Rabbit

### Isotype

IgG

### Purification


Affinity purification

### Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal, 50% glycerol, pH7.3.

## Validation Data



Western blot analysis of various lysates using OCIAD2 Rabbit pAb (A14962) at 1:1000 dilution.  
Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.  
Lysates/proteins: 25µg per lane.  
Blocking buffer: 3% nonfat dry milk in TBST.  
Detection: ECL Basic Kit (RM00020).  
Exposure time: 10s.



Immunofluorescence analysis of U2OS cells using OCIAD2 Rabbit pAb (A14962) at dilution of 1:100. Blue: DAPI for nuclear staining.