[KO Validated] Annexin A5/Annexin V Rabbit pAb

ABclomal[®]

Catalog No.: A1728 KO Validated

Basic Information

Observed MW 35kDa

Calculated MW 36kDa

Category Primary antibody

Applications ELISA,WB,IF/ICC

Cross-Reactivity Human, Mouse

Background

The Annexin 5 gene spans 29 kb containing 13 exons, and encodes a single transcript of approximately 1.6 kb and a protein product with a molecular weight of about 35 kDa.The protein encoded by this gene belongs to the annexin family of calcium-dependent phospholipid binding proteins some of which have been implicated in membrane-related events along exocytotic and endocytotic pathways. Annexin 5 is a phospholipase A2 and protein kinase C inhibitory protein with calcium channel activity and a potential role in cellular signal transduction, inflammation, growth and differentiation. Annexin 5 has also been described as placental anticoagulant protein I, vascular anticoagulant-alpha, endonexin II, lipocortin V, placental protein 4 and anchorin CII. Polymorphisms in this gene have been implicated in various obstetric complications.

Recommended Dilutions

Immunogen Information

WB	1:500 - 1:2000	Gene ID	Swiss Prot
IF/ICC	1:50 - 1:200	308	P08758

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 1-320 of human Annexin A5/Annexin V (NP_001145.1).

Synonyms

PP4; ANX5; ENX2; CPB-I; RPRGL3; HEL-S-7; VAC-alph; V

Contact


a 400-999-6126 x cn.market@abclonal.com.cn o www.abclonal.com.cn

Product Information

Source Rabbit **Isotype** IgG Purification Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.




Western blot analysis of extracts of various cell lines, using Annexin A5/Annexin V antibody (A1728) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.



Western blot analysis of extracts from normal (control) and Annexin A5/Annexin V knockout (KO) 293T cells, using Annexin A5/Annexin V antibody (A1728) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 1s.



Immunofluorescence analysis of HeLa cells using [KO Validated] Annexin A5/Annexin V Rabbit pAb (A1728) at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.