

SARS-CoV-2 Spike S1 mAb, BSA and glycerol free

Catalog No.: A20022 **6 Publications**

Basic Information

Observed MW

110kDa

Calculated MW**Category**

Primary antibody

Applications

ELISA, WB, FC, IF/ICC

Cross-Reactivity

SARS-CoV-2

CloneNo number

ARC2373

Background

Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) is an enveloped, positive-sense, single-stranded RNA virus that causes coronavirus disease 2019 (COVID-19). Virus particles include the RNA genetic material and structural proteins needed for invasion of host cells. Once inside the cell the infecting RNA is used to encode structural proteins that make up virus particles, nonstructural proteins that direct virus assembly, transcription, replication and host control and accessory proteins whose function has not been determined.~ The structural proteins of SARS-CoV-2 include the envelope protein (E), spike or surface glycoprotein (S), membrane protein (M) and the nucleocapsid protein (N). The spike glycoprotein is found on the outside of the virus particle and gives coronavirus viruses their crown-like appearance. This glycoprotein mediates attachment of the virus particle and entry into the host cell. S protein is an important target for vaccine development, antibody therapies and diagnostic antigen-based tests.

Recommended Dilutions

ELISA	1:1000 - 1:5000
WB	1:2000 - 1:10000
FC	1:50 - 1:200
IF/ICC	1:50 - 1:200

Immunogen Information

Gene ID

43740568

Swiss Prot**Immunogen**

Recombinant fusion protein of SARS-CoV-2 Spike S1.

Synonyms

spike glycoprotein

Contact

	400-999-6126
	cn.market@abclonal.com.cn
	www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage

Store at 4°C Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300, pH7.3.

Validation Data



Western blot analysis of lysates from control 293T and SARS-CoV-2 Spike S1-293T transfected cells, using SARS-CoV-2 Spike S1 mAb, BSA and glycerol free (A20022) at 1:10000 dilution.

Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Enhanced Kit (RM00021).

Exposure time: 180s.