# SARS-CoV-2 ORF9b Rabbit pAb

Catalog No.: A20260 1 Publications



## **Basic Information**

### **Observed MW**

12-14kDa

### **Calculated MW**

11kDa

#### Category

Primary antibody

## **Applications**

ELISA,WB

#### **Cross-Reactivity**

Human

# **Background**

SARS-CoV-2 (severe acute respiratory syndrome coronavirus 2) is the causative agent of the COVID19 pandemic.?? SARS-CoV-2 and SARS-CoVshare many proteins common in other CoVs, including 4 majorstructural proteins (S, E, M, and N proteins) and 16 nonstructural proteins (nsp1-16), they possess a unique set of proteins, namelyorf3a, 3b, 6, 7a, 7b, 8a, 8b, and 9b $\square$ The SARS-CoV-2 genome encodes for a small accessory protein termed Orf9b,?? which targets the mitochondrial outer membrane protein TOM70 in infected cells. This 98-amino acid (aa) protein is encoded by analternative open reading frame (ORF) within the N gene and istranslated via a leaky scanning mechanism during translationCrystal structures of orf9b alone revealed a homodimeric $\beta$ -strand-rich structure? with a hydro\_x005fphobic central tunnel for lipid binding, consistent with the role oforf9b in the mature virion assembly

## **Recommended Dilutions**

**WB** 

1:500 - 1:1000

# Immunogen Information

Gene ID

Swiss Prot

#### **Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 1-97 of coronavirus ORF9b (P0DTD2).

# **Synonyms**

## **Contact**

<b>a</b>	400-999-6126
×	cn.market@abclonal.com.cn
$\overline{\mathfrak{S}}$	www.abclonal.com.cn

## **Product Information**


SourceIsotypePurificationRabbitIgGAffinity purification

#### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.

# **Validation Data**



Western blot analysis of extracts of normal 293T cells 293T transfected with ORF9b Protein, using SARS-CoV-2 ORF9b Rabbit pAb (A20260) at 1:1000 dilution.

Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 30s.