

Transgelin (TAGLN) Rabbit mAb

Catalog No.: A21209

Recombinant

3 Publications

Basic Information

Observed MW

Refer to figures

Calculated MW

23kDa

Category

Primary antibody

Applications

ELISA, IHC-P, IF/ICC

Cross-Reactivity

Human, Mouse, Rat

CloneNo number

ARC53169

Background

This gene encodes a shape change and transformation sensitive actin-binding protein which belongs to the calponin family. It is ubiquitously expressed in vascular and visceral smooth muscle, and is an early marker of smooth muscle differentiation. The encoded protein is thought to be involved in calcium-independent smooth muscle contraction. It acts as a tumor suppressor, and the loss of its expression is an early event in cell transformation and the development of some tumors, coinciding with cellular plasticity. The encoded protein has a domain architecture consisting of an N-terminal calponin homology (CH) domain and a C-terminal calponin-like (CLIK) domain. Mice with a knockout of the orthologous gene are viable and fertile but their vascular smooth muscle cells exhibit alterations in the distribution of the actin filament and changes in cytoskeletal organization.

Recommended Dilutions

IHC-P 1:500 - 1:1000

IF/ICC 1:50 - 1:200

Immunogen Information

Gene ID

6876

Swiss Prot

Q01995

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 1-201 of human Transgelin (TAGLN) (NP_003177.2).

Synonyms

SM22; SMCC; TAGLN1; WS3-10; SM22-alpha; Transgelin (TAGLN)

Contact

 | 400-999-6126 | cn.market@abclonal.com.cn | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300, 0.05% BSA, 50% glycerol, pH7.3.


Validation Data




Immunofluorescence analysis of paraffin-embedded mouse brain using Transgelin (TAGLN) Rabbit mAb (A21209) at dilution of 1:100 (40x lens). Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.




Immunofluorescence analysis of paraffin-embedded mouse small intestine using Transgelin (TAGLN) Rabbit mAb (A21209) at dilution of 1:100 (40x lens). Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.




Confocal imaging of paraffin-embedded Human colon cancer using Transgelin (TAGLN) Rabbit mAb (A21209, dilution 1:200) followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). Objective: 40x. Perform high pressure antigen retrieval with 0.01 M citrate buffer (pH 6.0) prior to IF staining.




Confocal imaging of paraffin-embedded rat large intestine using Transgelin (TAGLN) Rabbit mAb (A21209, dilution 1:200) followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). Objective: 40x. Perform high pressure antigen retrieval with 0.01 M citrate buffer (pH 6.0) prior to IF staining.




Immunohistochemistry analysis of Transgelin (TAGLN) in paraffin-embedded human cervix cancer tissue using Transgelin (TAGLN) Rabbit mAb (A21209) at a dilution of 1:1000 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of Transgelin (TAGLN) in paraffin-embedded human colon tissue using Transgelin (TAGLN) Rabbit mAb (A21209) at a dilution of 1:1000 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of Transgelin (TAGLN) in paraffin-embedded mouse spleen tissue using Transgelin (TAGLN) Rabbit mAb (A21209) at a dilution of 1:1000 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0).



Immunohistochemistry analysis of Transgelin (TAGLN) in paraffin-embedded rat spleen tissue using Transgelin (TAGLN) Rabbit mAb (A21209) at a dilution of 1:1000 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0).

Validation Data

prior to IHC staining.

prior to IHC staining.