ABclonal www.abclonal.com

ABflo® 594 Rabbit anti-Human BTN3A1/2/3 mAb

Catalog No.: A24823

Basic Information

Observed MW

Calculated MW

58kDa

Category

Primary antibody

Applications

FC

Cross-Reactivity

Human

CloneNo number

ARC63604-ABflo594

Conjugate

ABflo® 594. Ex:588nm. Em:604nm.

Background

The butyrophilin (BTN) genes are a group of major histocompatibility complex (MHC)-associated genes that encode type I membrane proteins with 2 extracellular immunoglobulin (Ig) domains and an intracellular B30.2 (PRYSPRY) domain. Three subfamilies of human BTN genes are located in the MHC class I region: the single-copy BTN1A1 gene (MIM 601610) and the BTN2 (e.g., BTN2A1; MIM 613590) and BTN3 (e.g., BNT3A1) genes, which have undergone tandem duplication, resulting in 3 copies of each (summary by Smith et al., 2010 [PubMed 20208008]).

Recommended Dilutions

FC

5 μ l per 10^6 cells in 100 μ l volume

Immunogen Information

Gene ID

Swiss Prot

O00481/P78410/O00478

Immunogen

11119/11118/10384

Recombinant fusion protein containing a sequence corresponding to amino acids 470-630 of human BTN3A1(NP_008979.3).

Synonyms

BTF5; BT3.1; CD277; BTN3.1

Contact

2		400-999-6126
\bowtie	Τ	cn.market@abclonal.com.cn
•	Т	www.abclonal.com.cn

Product Information


SourceIsotypePurificationRabbitIgGAffinity purification


Storage

Store at 2-8°C. Avoid freeze.

Buffer: PBS with 0.03% proclin300,0.2% BSA,pH7.3.

Validation Data





Flow cytometry: 1X10^6 MCF7 cells (Low Expression,left) and U266 cells (right) were surface-stained with ABflo® 594 Rabbit anti-Human BTN3A1/2/3 mAb (A24823,5 μ I/Test,orange line) or ABflo® 594 Rabbit IgG isotype control (A23821,5 μ I/Test,blue line). Non-fluorescently stained cells were used as blank control (red line).

Flow cytometry: $1X10^6$ U266 cells were surface-stained with ABflo® 594 Rabbit IgG isotype control (A23821,5 μ I/Test,left) or ABflo® 594 Rabbit anti-Human BTN3A1/2/3 mAb (A24823,5 μ I/Test,right).