# ABflo® 488 Rabbit anti-Mouse CD215/IL-15Ra mAb

Catalog No.: A25573



## **Basic Information**

**Observed MW** Refer to figures

Calculated MW 28kDa

Category Primary antibody

Applications IF/ICC,FC

Cross-Reactivity Mouse

CloneNo number ARC66399-ABflo488

#### Conjugate

ABflo® 488. Ex:491nm. Em:516nm.

## **Recommended Dilutions**

IF/ICC	1:50 - 1:200
FC	5 μl per 10^6 cells in 100 μl volume

## Background

Predicted to enable interleukin-15 receptor activity and protein kinase binding activity. Acts upstream of or within positive regulation of natural killer cell differentiation. Located in cytoplasmic vesicle and plasma membrane. Is expressed in central nervous system; retina; retina inner nuclear layer; and retina outer nuclear layer. Orthologous to human IL15RA (interleukin 15 receptor subunit alpha).

# Immunogen Information

Gene ID 16169 Swiss Prot Q60819

#### Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 33-205 of mouse CD215/IL-15R $\alpha$  (NP\_032384.1).

#### Synonyms

ll15ra

## Contact

6		400-999-6126
$\times$		cn.market@abclonal.com.cn
€		www.abclonal.com.cn


# **Product Information**


**Source** Rabbit **Isotype** IgG **Purification** Affinity purification

#### Storage

Store at 2-8°C. Avoid freeze. Buffer: PBS with 0.03% proclin300,0.2% BSA,pH7.3.

## Validation Data





Flow cytometry: 1X10^6 CHO cells (negative control,left) and CHO (Transfection,right) cells were surface-stained with ABflo® 488 Rabbit anti-Mouse CD215/IL-15R $\alpha$  mAb (A25573,5 µl/Test,orange line) or ABflo® 488 Rabbit IgG isotype control (A22069,5 µl/Test,blue line). Non-fluorescently stained cells were used as blank control (red line).

Confocal imaging of 293F cells transfected with CD215/IL-15R $\alpha$  using ABflo® 488 Rabbit anti-Mouse CD215/IL-15R $\alpha$  mAb (A25573, dilution 1:50). DAPI was used for nuclear staining (Blue). Objective: 100x.