Leader in Biomolecular Solutions for Life Science

www.abclonal.com

CD45 Rabbit mAb

Catalog No.: A25762 Recombinant

Basic Information

Observed MW

Calculated MW
131kda/136kDa/141kDa/143kDa/147kDa

Category
Primary antibody

Applications
FC,ELISA

Cross-Reactivity
Human

CloneNo number
ARC5024

Background

The protein encoded by this gene is a member of the protein tyrosine phosphatase (PTP) family. PTPs are known to be signaling molecules that regulate a variety of cellular processes including cell growth, differentiation, mitosis, and oncogenic transformation. This PTP contains an extracellular domain, a single transmembrane segment and two tandem intracytoplasmic catalytic domains, and thus is classified as a receptor type PTP. This PTP has been shown to be an essential regulator of T- and B-cell antigen receptor signaling. It functions through either direct interaction with components of the antigen receptor complexes, or by activating various Src family kinases required for the antigen receptor signaling. This PTP also suppresses JAK kinases, and thus functions as a regulator of cytokine receptor signaling. Alternatively spliced transcripts variants of this gene, which encode distinct isoforms, have been reported.

Recommended Dilutions

FC	1:500 - 1:1000
ELISA	Recommended starting concentration is 1 μg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID
5788

Swiss Prot
P08575

Immunogen
Recombinant protein (or fragment).This information is considered to be commercially sensitive.

Synonyms
LCA; LY5; B220; CD45; L-CA; T200; CD45R; GP180; IMD105

Contact

400-999-6126

cn.market@abclonal.com.cn

www.abclonal.com.cn

Product Information


Source
Rabbit

Isotype
IgG


Purification
Affinity purification

Storage
Store at -20℃

Validation Data



Flow cytometry: 1X10^6 293T cells (negative control,left) and Jurkat cells (right) were surface-stained with CD45 Rabbit mAb (A25762,2 µg/mL,orange line) or Rabbit IgG isotype control (AC042,2 µg/mL,blue line), followed by ABflo® 405-conjugated goat anti-rabbit pAb staining. Non-fluorescently stained cells were used as blank control (red line).



Flow cytometry: 1X10^6 Jurkat cells were surface-stained with Rabbit IgG isotype control (AC042,2 µg/mL,left) or CD45 Rabbit mAb (A25762,2 µg/mL,right), followed by ABflo® 405-conjugated goat anti-rabbit pAb staining.