# **HNRNPR Rabbit pAb**

Catalog No.: A5883



### **Basic Information**

#### **Observed MW**

71kDa/77kDa/80kDa

### **Calculated MW**

71kDa

#### Category

Primary antibody

### **Applications**

ELISA,WB,IHC-P

#### **Cross-Reactivity**

Human, Mouse, Rat

### **Background**

This gene encodes an RNA-binding protein that is a member of the spliceosome C complex, which functions in pre-mRNA processing and transport. The encoded protein also promotes transcription at the c-fos gene. Alternative splicing results in multiple transcript variants. There are pseudogenes for this gene on chromosomes 4, 11, and 10.

### **Recommended Dilutions**

**WB** 1:500 - 1:2000

IHC-P 1:100 - 1:200

### **Immunogen Information**

**Gene ID Swiss Prot** 10236 043390

#### **Immunogen**

A synthetic peptide corresponding to a sequence within amino acids 500 to the C-terminus of human HNRNPR (NP\_005817.1).

### **Synonyms**

HNRPR; NEDDFSB; hnRNP-R; HNRNPR

### **Contact**

<b>a</b>		400-999-6126
$\bowtie$		cn.market@abclonal.com.cn
$\overline{a}$	ı	www.ahclonal.com.cn

### **Product Information**


SourceIsotypePurificationRabbitIgGAffinity purification

### **Storage**

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

## **Validation Data**



Western blot analysis of various lysates using HNRNPR Rabbit pAb (A5883) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates / proteins: 25  $\mu$ g per lane.

Blocking buffer: 3 % nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 1s.