Synaptophysin Rabbit pAb

Catalog No.: A6344 15 Publications



Basic Information

Observed MW

38kDa

Calculated MW

34kDa

Category

Primary antibody

Applications

ELISA, WB, IHC-P, IF/ICC, IP

Cross-Reactivity

Human, Mouse, Rat

Background

This gene encodes an integral membrane protein of small synaptic vesicles in brain and endocrine cells. The protein also binds cholesterol and is thought to direct targeting of vesicle-associated membrane protein 2 (synaptobrevin) to intracellular compartments. Mutations in this gene are associated with an X-linked form of cognitive disability.

Recommended Dilutions

WB 1:500 - 1:1000

IHC-P 1:50 - 1:200

IF/ICC 1:50 - 1:200

IP 0.5μg-4μg antibody for 400μg-600μg extracts of

whole cells

Immunogen Information

Gene ID6855
Swiss Prot
P08247

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 222-313 of human Synaptophysin (NP_003170.1).

Synonyms

MRX96; MRXSYP; XLID96; Synaptophysin

Contact

a	400-999-6126
×	cn.market@abclonal.com.cn
\overline{a}	www.abclonal.com.cn


Product Information

SourceIsotypePurificationRabbitIgGAffinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.




Western blot analysis of various lysates using Synaptophysin Rabbit pAb (A6344) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.


Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).


Exposure time: 1s.



Immunoprecipitation analysis of 600 μg extracts of Mouse brain cells using 3 μg Synaptophysin antibody (A6344). Western blot was performed from the immunoprecipitate using Synaptophysin antibody (A6344) at a dilution of 1:1000.




Immunohistochemistry analysis of paraffinembedded Human brain using Synaptophysin Rabbit pAb (A6344) at dilution of 1:200 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.




Immunohistochemistry analysis of paraffinembedded Mouse pancreatic islet using Synaptophysin Rabbit pAb (A6344) at dilution of 1:200 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.




Immunohistochemistry analysis of paraffinembedded Rat brain using Synaptophysin Rabbit pAb (A6344) at dilution of 1:200 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.




Immunofluorescence analysis of Neuro-2a cells using Synaptophysin Rabbit pAb (A6344) at dilution of 1:100 (40x lens). Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.




Immunofluorescence analysis of SH-SY5Y cells using Synaptophysin Rabbit pAb (A6344) at dilution of 1:100 (40x lens). Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of paraffinembedded rat brain using Synaptophysin Rabbit pAb (A6344) at dilution of 1:100 (40x lens). Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of paraffinembedded human pancreas using Synaptophysin Rabbit pAb (A6344) at dilution of 1:100 (40x lens). Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of paraffinembedded mouse brain using Synaptophysin Rabbit pAb (A6344) at dilution of 1:100 (40x lens). Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.