# **KDM4C Rabbit pAb**

Catalog No.: A8485 2 Publications



### **Basic Information**

#### **Observed MW**

120kDa

### **Calculated MW**

120kDa

### Category

Primary antibody

### **Applications**

ELISA,WB

### **Cross-Reactivity**

Human, Mouse, Rat

## **Background**

This gene is a member of the Jumonji domain 2 (JMJD2) family. The encoded protein is a trimethylation-specific demethylase, and converts specific trimethylated histone residues to the dimethylated form. This enzymatic action regulates gene expression and chromosome segregation. Chromosomal aberrations and changes in expression of this gene may be found in tumor cells. Alternative splicing results in multiple transcript variants.

## **Recommended Dilutions**

**WB** 

1:500 - 1:2000

## **Immunogen Information**

**Gene ID** 23081

**Swiss Prot** 

Q9H3R0

### **Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 640-830 of human KDM4C (NP\_055876.2).

## **Synonyms**

GASC1; JHDM3C; JMJD2C; TDRD14C; KDM4C

## **Contact**

6		400-999-6126
$\bowtie$		cn.market@abclonal.com.cn
•	T	www.abclonal.com.cn


## **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

### **Storage**

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.



Western blot analysis of Mouse brain, using KDM4C Rabbit pAb (A8485) at 1:800 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 60s.