# Mouse anti mCherry-Tag mAb

Catalog No.: AE002 20 Publications



# **Basic Information**

### **Observed MW**

55kDa

#### **Calculated MW**

### Category

Tag antibody

# **Applications**

ELISA,WB

### **Cross-Reactivity**

Species independent

### CloneNo number

AMC0502

# **Background**

Protein tags are peptide sequences genetically grafted onto a recombinant protein. Often these tags are removable by chemical agents or by enzymatic means, such as proteolysis or intein splicing. Tags are attached to proteins for various purposes. Epitope tags are short peptide sequences which are chosen because high-affinity antibodies can be reliably produced in many different species. These are usually derived from viral genes, which explain their high immunoreactivity. Epitope tags include V5-tag, Myc-tag, HA-tag and NE-tag. These tags are particularly useful for western blotting, immunofluorescence and immunoprecipitation experiments, although they also find use in antibody purification.

**Swiss Prot** 

### **Recommended Dilutions**

**WB** 

1:5000 - 1:10000

# **Immunogen Information**

Gene ID

#### **Immunogen**

Recombinant protein of mCherry tag

#### **Synonyms**

mCherry;mCherry tag;mCherry-tag

### **Contact**

<b>a</b>		400-999-6126
$\bowtie$		cn.market@abclonal.com.cn
$\odot$	T	www.abclonal.com.cn


### **Product Information**

SourceIsotypePurificationMouseIqG1Affinity purification

#### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,1% BSA,50% glycerol,pH7.3.



Western blot analysis of extracts of normal 293T cells and 293T transfected with CT55 Protein, using Mouse anti mCherry-Tag antibody (AE002) at 1:5000 dilution.

Secondary antibody: HRP Goat Anti-Mouse IgG (H+L) (AS003) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 1s.