

Mouse anti Myc-Tag mAb

Catalog No.: AE010 **96 Publications**

Basic Information

Observed MW

50-70kDa/58kDa

Calculated MW**Category**

Tag antibody

Applications

ELISA, WB, IF, IP

Cross-Reactivity

Species independent

CloneNo number

AMC0504

Background

Protein tags are peptide sequences genetically grafted onto a recombinant protein. Often these tags are removable by chemical agents or by enzymatic means, such as proteolysis or intein splicing. Tags are attached to proteins for various purposes. Epitope tags are short peptide sequences which are chosen because high-affinity antibodies can be reliably produced in many different species. These are usually derived from viral genes, which explain their high immunoreactivity. Epitope tags include V5-tag, Myc-tag, HA-tag and NE-tag. These tags are particularly useful for western blotting, immunofluorescence and immunoprecipitation experiments, although they also find use in antibody purification.

Recommended Dilutions

WB 1:2000 - 1:6000**IF** 1:1000-1:5000**IP** 0.5µg-4µg antibody for
200µg-400µg extracts of
whole cells

Immunogen Information

Gene ID**Swiss Prot****Immunogen**

A synthetic peptide corresponding to Myc tag.

Synonyms

Myc;Myc tag;Myc-tag

Contact

 | 400-999-6126 | cn.market@abclonal.com.cn | www.abclonal.com.cn

Product Information

Source

Mouse

Isotype

IgG1

Purification


Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal, 50% glycerol, pH7.3.

Validation Data



Western blot analysis of extracts of normal 293T cells and 293T transfected with Myc-NLK protein, using Mouse anti Myc-Tag mAb (AE010) at 1:5000 dilution.


Secondary antibody: HRP Goat Anti-Mouse IgG (H+L) (AS003) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.


Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 1s.



Immunoprecipitation of NLK-Myc from 300 µg extracts of 293F cells transfected with a NLK expression vector containing a single C-terminal Myc-Tag was performed using 3 µg of Mouse anti Myc-Tag mAb (AE010). Mouse IgG isotype control (AC011) was used to precipitate the Control IgG sample. IP samples were eluted with 1X Laemmli Buffer. The Input lane represents 10 % of the total input. Western blot analysis of immunoprecipitates was conducted using Mouse anti Myc-Tag mAb (AE102) at a dilution of 1:3000.



Confocal imaging of 293F cells transfected with SQSTM1-Myc(N) cells using Mouse anti Myc-Tag mAb (AE010, dilution 1:2000) followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). Objective: 100x.