

Affinity Gel-conjugated Rabbit anti DDDDK-Tag mAb mAb {Anti-Flag □□□□}

Catalog No.: AE121

Basic Information

Observed MW

40kDa

Calculated MW

Category

Tag antibody

Applications

IP

Cross-Reactivity

Species independent

Conjugate

Affinity Gel

Background

FLAG-tag, or FLAG octapeptide, or FLAG epitope, is a polypeptide protein tag that can be added to a protein using recombinant DNA technology, having the sequence motif DYKDDDDK. It has been used for studying proteins in living cells and for protein purification by affinity chromatography. It has been used to separate recombinant, overexpressed protein from wild-type protein expressed by the host organism. It can also be used in the isolation of protein complexes with multiple subunits, because its mild purification procedure tends not to disrupt such complexes. It has been used to obtain proteins of sufficient purity and quality to carry out 3D structure determination by x-ray crystallography. A FLAG-tag can be used in many different assays that require recognition by an antibody. If there is no antibody against a given protein, adding a FLAG-tag to a protein allows the protein to be studied with an antibody against the FLAG sequence. Examples are cellular localization studies by immunofluorescence or detection by SDS PAGE protein electrophoresis and Western blotting.

Recommended Dilutions

IP 20μl-40μl Affinity Gel for
100μg-300μg extracts of
whole cells

Immunogen Information

Gene ID

Swiss Prot

Immunogen

A synthetic peptide corresponding to DDDDK tag.

Synonyms

DDDDK; DDDDK tag; DDDDK-tag

Contact

☎ | 400-999-6126

✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification

Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300, 0.05% BSA, 50% glycerol, pH7.3.

Validation Data

Immunoprecipitation of DDDDK-Tag in 150 µg extracts from 293F cells transfected with DDDDK-Tag using 20 µl Affinity Gel-conjugated Rabbit anti DDDDK-Tag mAb (AE121). Western blot analysis was performed using Affinity Gel-conjugated Rabbit anti DDDDK-Tag mAb (AE024) at 1:5000 dilution.

