

# Phospho-Smad2-S465/S467 + Smad3-S423/S425 Rabbit mAb

Catalog No.: AP1343 Recombinant 1 Publications

## **Basic Information**

## **Observed MW**

52kDa/60kDa

#### **Calculated MW**

48kDa/52kDa/25kDa/35kDa/43kDa

#### Category

Primary antibody

#### **Applications**

ELISA,WB

#### **Cross-Reactivity**

Human, Mouse, Rat

#### CloneNo number

ARC56314

# **Background**

The protein encoded by this gene belongs to the SMAD, a family of proteins similar to the gene products of the Drosophila gene 'mothers against decapentaplegic' (Mad) and the C. elegans gene Sma. SMAD proteins are signal transducers and transcriptional modulators that mediate multiple signaling pathways. This protein mediates the signal of the transforming growth factor (TGF)-beta, and thus regulates multiple cellular processes, such as cell proliferation, apoptosis, and differentiation. This protein is recruited to the TGF-beta receptors through its interaction with the SMAD anchor for receptor activation (SARA) protein. In response to TGF-beta signal, this protein is phosphorylated by the TGF-beta receptors. The phosphorylation induces the dissociation of this protein with SARA and the association with the family member SMAD4. The association with SMAD4 is important for the translocation of this protein into the nucleus, where it binds to target promoters and forms a transcription repressor complex with other cofactors. This protein can also be phosphorylated by activin type 1 receptor kinase, and mediates the signal from the activin. Alternatively spliced transcript variants have been observed for this gene. [provided by RefSeq, May 2012]

## **Recommended Dilutions**

**WB** 

1:1000 - 1:5000

## Immunogen Information

**Gene ID** 4087/4088

**Swiss Prot** 

38 Q15796/P84022

#### **Immunogen**

A synthetic phosphorylated peptide around S465 & S467 of human SMAD2 (NP\_005892.1).

# **Synonyms**

 $HSPC193; \ HsT17436; \ JV15-2; \ LDS1C; \ LDS3; \ MADH3; \ Phospho-Smad2-S465/S467 + Smad3-S423/S425$ 

## **Contact**

<b>a</b>	400-999-6126
$\sim$	cn.market@abclonal.com.cn
<b>⊙</b>	www.abclonal.com.cn

### **Product Information**


SourceIsotypePurificationRabbitIgGAffinity purification

#### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300,0.05% BSA,50% glycerol,pH7.3.

## **Validation Data**



Western blot analysis of various lysates, using Phospho-Smad2-S465/S467 + Smad3-S423/S425 Rabbit mAb (AP1343) at1:2000 dilution.HeLa and NIH/3T3 cells were treated by TGF- $\beta3$ [H[] (200 ng/mL) at 37°C for 30 minutes after serum-starvation overnight.


Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 60s.



Western blot analysis of various lysates using Phospho-Smad2-S465/S467 + Smad3-S423/S425 Rabbit mAb (AP1343) at 1:1000 dilution.C6 cells were treated by TGF- $\beta$ 3 (200 ng/mL) at 37°C for 30 minutes after serum-starvation overnight.

Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 30s.