

FOXO1 Knockdown HeLa Cell Line, Heterozygous

Catalog No.: RM01924

Basic Information

Catalog No.

RM01924

Category

Cell Line

Parental Cell line

HeLa

Genotype

Knockdown

Gene Information

Gene Symbol

FOXO1

Species

Human

Gene ID

2308

Swiss Prot

Q12778

Synonyms

FKH1; FKHR; FOXO1A

Contact

☎ | 400-999-6126

✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Background

This gene belongs to the forkhead family of transcription factors which are characterized by a distinct forkhead domain. The specific function of this gene has not yet been determined; however, it may play a role in myogenic growth and differentiation. Translocation of this gene with PAX3 has been associated with alveolar rhabdomyosarcoma. [provided by RefSeq, Jul 2008]

Product Information

Description

FOXO1 Knockdown HeLa Cell Line is engineered from HeLa cell line with Gene-Editing Technology.

Allele-1:1bp deletion in exon1

Allele-2:12bp deletion in exon1

Mammalian cells such as human, rat and mouse cells are normally diploid with two alleles.

Homozygote: both alleles were knocked out, mRNA has no signal, no expression of proteins.

Heterozygote: only one allele was knocked out, the mRNA transcript levels was decreased compared to wild type, and the protein expression levels was also lower than that of the wild type.

Packaging

1 vial parental cell line and 1 vial knockout cell line

Shipping Conditions

Dry ice

Amount

1~5x10⁶ cells/vial

Storage

Stored in liquid nitrogen for a long time less than -130°C. Minimizing freeze-thaw cycles.

Protocol

Upon arrival, it should be maintained in DMEM medium with 10%(v/v) fetal bovine serum and 100U penicillin-streptomycin, at 37°C with 5% CO₂ condition.

1. Thaw the vial in 37°C water bath, and shake it to melt as soon as possible.
2. Transfer the cell suspension to a 15mL conical tube with pre-warmed 5mL complete medium and centrifuge 1000rpm for approximately 5 minutes at room temperature.
3. Remove and discard the supernatant.
4. Resuspend the cell pellet with 1mL pre-warmed complete medium and seed in 10cm dish.
5. Add 8-10mL of complete medium.
6. Incubate the culture at 37°C incubator with 5% CO₂.
7. A subcultivation ratio of 1:2-1:4 is recommended.

Sequencing data

WT CGGCCCGCGCGGCCACCGGGGGCTGTGCGGGGAC
Mut CGGCCCGCGCGGCCA-CGGGGGGCTGTGCGGGGAC
Allele-1: 1bp deletion in exon1
WT GCGCGGCGCGCGG*****GGGGGCTGTGCGGG
Mut GCGCGGCGCGCGG***Deletion***GGGGGCTGTGCGGG
Allele-2: 12bp deletion in exon1

Genome sequence analysis of PCR products from parental (WT) and FOXO1 Knockdown (KD) HeLa cells, using sanger sequencing.