

Recombinant Human VEGFR-3/FLT-4 Protein

Catalog No.: RP00123 Recombinant

Sequence Information

Species Gene ID Swiss Prot Human 2324 P35916

Tags

C-hFc&His

Synonyms

FLT4;FLT-4;FLT41;LMPH1A;PCL;VEGFR-3; VEGFR3

Product Information

Source Purification HEK293 cells > 95% by SDS-PAGE.

Endotoxin

< 0.1 EU/ μ g of the protein by LAL method.

Formulation

Lyophilized from a 0.22 µm filtered solution of PBS, pH 7.4.Contact us for customized product form or formulation.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid votex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact

6	400-999-6126
\bowtie	cn.market@abclonal.com.cn
•	www.abclonal.com.cn

Background

The protein is a tyrosine kinase receptor for vascular endothelial growth factors C and D. The protein is thought to be involved in lymphangiogenesis and maintenance of the lymphatic endothelium. Mutations in this gene cause hereditary lymphedema type IA.

Basic Information

Description

Recombinant Human VEGFR-3/FLT-4 Protein is produced by HEK293 expression system. The target protein is expressed with sequence (Tyr25-Ile776) of human VEGFR3/FLT4 (Accession #NP_891555.2) fused with an Fc, 6×His tag at the C-terminus.

Bio-Activity

Measured by its binding ability in a functional ELISA. Immobilized Recombinant Human VEGF-C at 0.5 μ g/mL (100 μ L/well) can bind Recombinant Human VEGFR3 with a linear range of 3.92-15.70 ng/mL.


Storage

Store the lyophilized protein at -20 $^{\circ}\text{C}$ to -80 $^{\circ}\text{C}$ for long term.


After reconstitution, the protein solution is stable at -20 $^{\circ}$ C for 3 months, at 2-8 $^{\circ}$ C for up to 1 week.

Avoid repeated freeze/thaw cycles.

Validation Data



Recombinant Human VEGFR-3/FLT-4 Protein was determined by SDS-PAGE with Coomassie Blue, showing bands at 150-160 kDa, 100 kDa and 80 kDa.



Immobilized Recombinant Human VEGF-C, His Tag at 0.5 μ g/mL (100 μ L/well) can bind Recombinant Human VEGFR3,Fc Tag with a linear range of 3.92-15.70ng/mL.