

Recombinant Human Angiopoietin-2/ANGPT2 Protein

Catalog No.: RP02008

Recombinant

Sequence Information

Species	Gene ID	Swiss Prot
Human	285	O15123-1

Tags

C-His

Synonyms

AGPT2; ANG2;ANGPT2;ANG2

Product Information

Source	Purification
HEK293 cells	> 95% by SDS-PAGE.

Endotoxin

< 0.1 EU/μg of the protein by LAL method.

Formulation

Lyophilized from a 0.22 μm filtered solution of 20mM MOPS 150 mM NaCl 0.05% CHAPS pH 7.0.

Reconstitution

Centrifuge the tube before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Background

Angiopoietin-2 (Ang-2; also ANGPT2) is a secreted glycoprotein that plays a complex role in angiogenesis and inflammation. Mature Ang-2 is 478 amino acids in length. Ang2 is widely expressed during development, but it is restricted postnatally to highly angiogenic tissues such as the placenta, ovaries, and uterus. It is particularly abundant in vascular endothelial cells (EC) where it is stored in intracellular Weibel Palade bodies.

Basic Information

Description

Recombinant Human GTR Protein is produced by mammalian expression system. The target protein is expressed with sequence (Lys275-Phe496) of human Angiopoietin-2/ANGPT2 (Accession #O15123) fused with 6xHis at the C-terminus.

Bio-Activity

Storage

Store the lyophilized protein at -20°C to -80 °C for long term.

After reconstitution, the protein solution is stable at -20 °C for 3 months, at 2-8 °C for up to 1 week.

Avoid repeated freeze/thaw cycles.


Contact

☎ | 400-999-6126

✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Validation Data



Recombinant Human Angiopoietin-2/ANGPT2
Protein was determined by SDS-PAGE with
Coomassie Blue, showing a band at 35-45
kDa.