

Recombinant Human CXCL13/BLC Protein

Catalog No.: RP02839 **Recombinant**

Sequence Information

Species	Gene ID	Swiss Prot
Human	10563	O43927

Tags

No tag

Synonyms

ANGIE; ANGIE2; BCA-1; BCA1; BLC;
BLR1L; SCYB13; CXCL13

Product Information

Source	Purification
<i>E. coli</i>	> 95% by SDS-PAGE.

Endotoxin

<1EU/μg

Formulation

Lyophilized from a 0.22 μm filtered solution of 50 mM Tris, 0.5 M NaCl, pH 7.5.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Background

The chemokine CXCL13, also known as BCA-1 (B-cell-attracting chemokine-1) or BLC (B-lymphocyte chemoattractant), which belongs to the CXC chemokine family. CXCL13 and its receptor CXCR5 control the organization of B cells within follicles of lymphoid tissues. CXCL13 is known to dictate homing and motility of B cells in lymphoid tissue and has been implicated in the formation of ectopic lymphoid tissue in chronic inflammation. It involves in B-cell compartmental homing within secondary lymphoid organs and recently implicated in the pathogenesis of inflammatory and malignant lymphocyte-mediated diseases. In Primary central nervous system lymphoma (PCNSL), expression of BCA-1 by malignant lymphocytes and vascular endothelium may influence tumor development and localization to central nervous system (CNS). In T-lymphocytes, CXCL13 expression is thought to reflect a germinal center origin of the T-cell. CXCL13 expression may also provide an additional useful tool for the diagnosis of Angioimmunoblastic T-cell lymphoma (AITL).

Basic Information

Description

Recombinant Human CXCL13/BLC Protein is produced by *E. coli* expression system. The target protein is expressed with sequence (Val23-Arg94) of human CXCL13/BLC (Accession #NP_006410.1) fused with no additional amino acid.

Bio-Activity


Storage

Store the lyophilized protein at -20°C to -80°C for 12 months. After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week. Avoid repeated freeze/thaw cycles.

Contact

 | 400-999-6126 | cn.market@abclonal.com.cn | www.abclonal.com.cn

Validation Data



Recombinant Human CXCL13/BLC Protein was determined by SDS-PAGE with Coomassie Blue, showing a band at 10 kDa.