Leader in Biomolecular Solutions for Life Science

www.abclonal.com

[KO Validated] YAP1 Rabbit mAb

Catalog No.: A21216 KO Validated Recombinant 12 Publications

Basic Information

Observed MW
70 kDa

Calculated MW
54 kDa

Category
Primary antibody

Applications
WB,IP,IF/ICC,IF-P,IHC-P,ChIP,ELISA

Cross-Reactivity
Human, Mouse, Rat

CloneNo number
ARC53479

Background

This gene encodes a downstream nuclear effector of the Hippo signaling pathway which is involved in development, growth, repair, and homeostasis. This gene is known to play a role in the development and progression of multiple cancers as a transcriptional regulator of this signaling pathway and may function as a potential target for cancer treatment. Alternative splicing results in multiple transcript variants encoding different isoforms.

Recommended Dilutions

WB	1:20000 - 1:80000
IP	0.5μg-4μg antibody for 200μg-400μg extracts of whole cells
IF/ICC	1:200 - 1:800
IF-P	1:200 - 1:400
IHC-P	1:200 - 1:800
ChIP	5 μg antibody for 10 μg - 15 μg of Chromatin
ELISA	Recommended starting concentration is 1 μg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID
10413

Swiss Prot
P46937

Immunogen
Recombinant protein (or fragment).This information is considered to be commercially sensitive.

Synonyms
YAP; YKI; COB1; YAP2; YAP-1; YAP65; P1

Product Information

Source
Rabbit

Isotype
IgG

Purification
Affinity purification

Storage
Store at -20℃

Contact

☎	400-999-6126
✉	cn.market@abclonal.com.cn
🌐	www.abclonal.com.cn

Validation Data



Immunoprecipitation analysis of 300 µg extracts of HeLa cells using 3 µg YAP1 antibody (A21216).
Western blot was performed from the immunoprecipitate using [KO Validated] YAP1 Rabbit mAb (A21216) at a dilution of1:2000.



Western blot analysis of lysates from wild type (WT) and YAP1 knockout (KO) HeLa cells using [KO Validated] YAP1 Rabbit mAb (A21216) at 1:20000 dilution incubated overnight at 4℃.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25 µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 30s.



Western blot analysis of lysates from NIH/3T3 cells using [KO Validated] YAP1 Rabbit mAb (A21216) at 1:20000 dilution incubated overnight at 4℃.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25 µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 30s.

Validation Data



Chromatin immunoprecipitation analysis of extracts of HeLa cells, using 5 µg of [KO Validated] YAP1 Rabbit mAb (A21216) and rabbit IgG.The amount of immunoprecipitated DNA was checked by quantitative PCR. Histogram was constructed by the ratios of the immunoprecipitated DNA to the input.



Confocal imaging of C2C12 cells using [KO Validated] YAP1 Rabbit mAb (A21216, dilution 1:200) followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). The cells were counterstained with α-Tubulin Mouse mAb (AC012, dilution 1:400) followed by incubation with ABflo® 488-conjugated Goat Anti-Mouse IgG (H+L) Ab (AS076, dilution 1:500) (Green). DAPI was used for nuclear staining (Blue). Objective: 100x.



Confocal imaging of HeLa cells using [KO Validated] YAP1 Rabbit mAb (A21216, dilution 1:200) followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). The cells were counterstained with α-Tubulin Mouse mAb (AC012, dilution 1:400) followed by incubation with ABflo® 488-conjugated Goat Anti-Mouse IgG (H+L) Ab (AS076, dilution 1:500) (Green). DAPI was used for nuclear staining (Blue). Objective: 100x.



Confocal imaging of paraffin-embedded Human colon tissue using [KO Validated] YAP1 Rabbit mAb (A21216, dilution 1:400) followed by a further incubation with Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IF staining. Objective: 40x.



Confocal imaging of paraffin-embedded Mouse intestine tissue using [KO Validated] YAP1 Rabbit mAb (A21216, dilution 1:400) followed by a further incubation with Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IF staining. Objective: 40x.



Confocal imaging of paraffin-embedded Mouse testis tissue using [KO Validated] YAP1 Rabbit mAb (A21216, dilution 1:400) followed by a further incubation with Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IF staining. Objective: 40x.



Confocal imaging of paraffin-embedded Rat intestine tissue using [KO Validated] YAP1 Rabbit mAb (A21216, dilution 1:400) followed by a further incubation with Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IF staining. Objective: 40x.

Validation Data



Immunohistochemistry analysis of paraffin-embedded human breast cancer tissue using [KO Validated] YAP1 Rabbit mAb (A21216) at a dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded human thyroid cancer tissue using [KO Validated] YAP1 Rabbit mAb (A21216) at a dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded human tonsil tissue using [KO Validated] YAP1 Rabbit mAb (A21216) at a dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded mouse spleen tissue using [KO Validated] YAP1 Rabbit mAb (A21216) at a dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded rat spleen tissue using [KO Validated] YAP1 Rabbit mAb (A21216) at a dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IHC staining.