Leader in Biomolecular Solutions for Life Science

www.abclonal.com

EIF4G2 Knockdown A549 Cell Lysate, Heterozygous

Catalog No.: RM50037

Basic Information

Catalog No.
RM50037

Category
Cell Lysate

Parental Cell line
A549

Genotype
Knockdown

Gene Information

Gene Symbol
EIF4G2

Species
Human

Gene ID
1982

Swiss Prot
P78344

Synonyms
P97; AAG1; DAP5; NAT1; EIF4G2/p97

Contact

400-999-6126

cn.market@abclonal.com.cn

www.abclonal.com.cn

Background

Translation initiation is mediated by specific recognition of the cap structure by eukaryotic translation initiation factor 4F (eIF4F), which is a cap binding protein complex that consists of three subunits: eIF4A, eIF4E and eIF4G. The protein encoded by this gene shares similarity with the C-terminal region of eIF4G that contains the binding sites for eIF4A and eIF3; eIF4G, in addition, contains a binding site for eIF4E at the N-terminus. Unlike eIF4G, which supports cap-dependent and independent translation, this gene product functions as a general repressor of translation by forming translationally inactive complexes. In vitro and in vivo studies indicate that translation of this mRNA initiates exclusively at a non-AUG (GUG) codon. Alternatively spliced transcript variants encoding different isoforms of this gene have been described.

Product Information

Description
EIF4G2 Knockdown A549 cell line is engineered from A549 cell line with Gene-Editing Technology.
Allele-1:exon1 was deleted
Allele-2:WT
Mammalian cells such as human, rat and mouse cells are normally diploid with two alleles. Homozygote: both alleles were knocked out, mRNA has no signal, no expression of proteins. Heterozygote: only one allele was knocked out, the mRNA transcript levels was decreased compared to wild type, and the protein expression levels was also lower than that of the wild type.

Packaging
1 vial parental cell Lysate and 1 vial knockout cell Lysate

Shipping Conditions
4℃

Amount
50μL, 2μg/μL.

Storage
Lysate is stable for 12 months when stored at -20℃. Minimizing freeze-thaw cycles.

Protocol
To be used as WB control. Lysate is supplied in 1× SDS sample buffer (2% SDS, 60 mM Tris-HCl pH 6.8, 10% Glycerol, 0.02% Bromophenol blue, 60 mM beta-mercaptoethanol). Lysate should be boiled for 3 - 5 minutes before loading onto gel.

Sequencing data



Genome sequence analysis of PCR products from parental (WT) and EIF4G2 knockdown (KD) A549 cells, using sanger sequencing.