

Catalog No.: A0009 2 Publications



## **Basic Information**

Observed MW 110kDa

Calculated MW 93kDa

Category Primary antibody

Applications WB

Cross-Reactivity Human

# Background

This gene encodes the alpha subunit of transcription factor hypoxia-inducible factor-1 (HIF-1), which is a heterodimer composed of an alpha and a beta subunit. HIF-1 functions as a master regulator of cellular and systemic homeostatic response to hypoxia by activating transcription of many genes, including those involved in energy metabolism, angiogenesis, apoptosis, and other genes whose protein products increase oxygen delivery or facilitate metabolic adaptation to hypoxia. HIF-1 thus plays an essential role in embryonic vascularization, tumor angiogenesis and pathophysiology of ischemic disease. Alternatively spliced transcript variants encoding different isoforms have been identified for this gene.

### **Recommended Dilutions**

1:200 - 1:2000

# **Immunogen Information**

WB

#### Gene ID 3091

Swiss Prot Q16665

#### Immunogen

Recombinant protein of Human HIF1 $\alpha$ 

**Product Information** 

#### Synonyms

HIF1; MOP1; PASD8; HIF-1A; bHLHe78; HIF-1alpha; HIF1-ALPHA; HIF-1-alpha; HIF1 $\alpha$ 

Contact	
---------	--

6

 $\sim$ 

€

# 400-999-6126 Source


cn.market@abclonal.com.cn

www.abclonal.com.cn

**lsotype** IgG Purification Affinity purification

#### Storage

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.



Western blot analysis of lysates from HeLa cells, using HIF1 $\alpha$  Rabbit pAb (A0009). Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST.