VEGFR1 Rabbit pAb

Catalog No.: A0058 1 Publications



Basic Information

Observed MW

200kDa

Calculated MW

151kDa

Category

Primary antibody

Applications

WB,IF/ICC,ELISA

Cross-Reactivity

Human, Mouse, Rat

Background

This gene encodes a member of the vascular endothelial growth factor receptor (VEGFR) family. VEGFR family members are receptor tyrosine kinases (RTKs) which contain an extracellular ligand-binding region with seven immunoglobulin (Ig)-like domains, a transmembrane segment, and a tyrosine kinase (TK) domain within the cytoplasmic domain. This protein binds to VEGFR-A, VEGFR-B and placental growth factor and plays an important role in angiogenesis and vasculogenesis. Expression of this receptor is found in vascular endothelial cells, placental trophoblast cells and peripheral blood monocytes. Multiple transcript variants encoding different isoforms have been found for this gene. Isoforms include a full-length transmembrane receptor isoform and shortened, soluble isoforms. The soluble isoforms are associated with the onset of pre-eclampsia.

Recommended Dilutions

WB 1:500 - 1:1000

IF/ICC 1:50 - 1:200

 $\begin{array}{c} \textbf{ELISA} & \text{Recommended starting} \\ & \text{concentration is 1 } \mu\text{g/mL}. \end{array}$

Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID2321

Swiss Prot
P17948

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

FLT; FLT-1; VEGFR1; VEGFR-1

Contact

<u>a</u>		400-999-6126
\bowtie		cn.market@abclonal.com.cn
•	Ī	www.abclonal.com.cn


Product Information

SourceIsotypePurificationRabbitIgGAffinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.


Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.




Western blot analysis of various lysates using VEGFR1 Rabbit pAb (A0058) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020).

Exposure time: 180s.



Immunofluorescence analysis of NIH/3T3 cells using VEGFR1 Rabbit pAb (A0058) at dilution of 1:100. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of U2OS cells using VEGFR1 Rabbit pAb (A0058) at dilution of 1:100. Blue: DAPI for nuclear staining.