Leader in Biomolecular Solutions for Life Science

www.abclonal.com

c-Fos Rabbit pAb

Catalog No.: A0236SP **33 Publications**

Basic Information

Observed MW
62 kDa

Calculated MW
41 kDa/36 kDa/29 kDa

Category
Primary antibody

Applications
WB,IF-P,IHC-P,ELISA

Cross-Reactivity
Human, Mouse, Rat

Background

The Fos gene family consists of 4 members: FOS, FOSB, FOSL1, and FOSL2. These genes encode leucine zipper proteins that can dimerize with proteins of the JUN family, thereby forming the transcription factor complex AP-1. As such, the FOS proteins have been implicated as regulators of cell proliferation, differentiation, and transformation. In some cases, expression of the FOS gene has also been associated with apoptotic cell death.

Recommended Dilutions

WB	1:1000 - 1:2000
IF-P	1:200 - 1:400
IHC-P	1:200 - 1:800
ELISA	Recommended starting concentration is 1 μg/mL. Please optimize the concentration based on your specific assay requirements.For high-ratio antibody dilutions (≥1:10000)

Immunogen Information

Gene ID
2353

Swiss Prot
P01100

Immunogen
This information is considered to be commercially sensitive.

Synonyms
p55; AP-1; C-FOS; c-Fos

Contact

400-999-6126

cn.market@abclonal.com.cn

www.abclonal.com.cn

Product Information

Source
Rabbit

Isotype
IgG

Purification
Affinity purification

Storage
Store at -20℃

Validation Data



Western blot analysis of lysates from HeLa cells using c-Fos Rabbit pAb (A0236SP) at 1:1000 dilution incubated overnight at 4℃. HeLa cells were treated with TPA (200 nM) at 37°C for 4 hours.|Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.|Lysates/proteins: 30 µg per lane.| Blocking buffer: 3% nonfat dry milk in TBST.| Detection: ECL Basic Kit (RM00020).| Exposure time: 45 s



Confocal imaging of paraffin-embedded Human brain tissue using c-Fos Rabbit pAb (A0236SP, dilution 1:200) followed by a further incubation with Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IF staining. Objective: 40x



Confocal imaging of paraffin-embedded Mouse brain tissue using c-Fos Rabbit pAb (A0236SP, dilution 1:200) followed by a further incubation with Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IF staining. Objective: 40x



Confocal imaging of paraffin-embedded Rat brain tissue using c-Fos Rabbit pAb (A0236SP, dilution 1:200) followed by a further incubation with Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IF staining. Objective: 40x



Immunohistochemistry analysis of paraffin-embedded Human colon tissue using c-Fos Rabbit pAb (A0236SP) at a dilution of 1:400 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining



Immunohistochemistry analysis of paraffin-embedded Human brain tissue using c-Fos Rabbit pAb (A0236SP) at a dilution of 1:400 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining



Immunohistochemistry analysis of paraffin-embedded Human tonsil tissue using c-Fos Rabbit pAb (A0236SP) at a dilution of 1:400 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining

Validation Data



Immunohistochemistry analysis of paraffin-embedded Mouse colon tissue using c-Fos Rabbit pAb (A0236SP) at a dilution of 1:400 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining



Immunohistochemistry analysis of paraffin-embedded Rat brain tissue using c-Fos Rabbit pAb (A0236SP) at a dilution of 1:400 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining



Immunohistochemistry analysis of paraffin-embedded Rat colon tissue using c-Fos Rabbit pAb (A0236SP) at a dilution of 1:400 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining

For research use only. Not for therapeutic or diagnostic purposes.
Please visit http://abclonal.com for a complete listing of recommended products.