

PPAR γ Rabbit pAb

Catalog No.: A0270SP

78 Publications

Basic Information

Observed MW

57 kDa

Calculated MW

58 kDa/55 kDa/22 kDa

Category

Primary antibody

Applications

WB, IF/ICC, IHC-P, ELISA

Cross-Reactivity

Human, Mouse, Rat

CloneNo number

Unconjugated

Background

This gene encodes a member of the peroxisome proliferator-activated receptor (PPAR) subfamily of nuclear receptors. PPARs form heterodimers with retinoid X receptors (RXRs) and these heterodimers regulate transcription of various genes. Three subtypes of PPARs are known: PPAR-alpha, PPAR-delta, and PPAR-gamma. The protein encoded by this gene is PPAR-gamma and is a regulator of adipocyte differentiation. Additionally, PPAR-gamma has been implicated in the pathology of numerous diseases including obesity, diabetes, atherosclerosis and cancer. Alternatively spliced transcript variants that encode different isoforms have been described.

Recommended Dilutions

WB 1:2000 - 1:5000**IF/ICC** 1:200 - 1:2000**IHC-P** 1:500 - 1:2000

ELISA Recommended starting concentration is 1 μ g/mL. Please optimize the concentration based on your specific assay requirements. For high-ratio antibody dilutions ($\geq 1:10000$) a sequential dilution method is strongly recommended to ensure measurement accuracy.

Contact

 | 400-999-6126 | cn.market@abclonal.com.cn

Immunogen Information

Gene ID

5468

Swiss Prot

P37231

Immunogen

This information is considered to be commercially sensitive.

Synonyms

GLM1; CIMT1; NR1C3; PPARG1; PPARG2; PPARG5; PPARGgamma

Product Information

Source

Rabbit

Isotype

IgG

Purification

Affinity purification



Storage

Store at -20°C




www.abclonal.com.cn


Validation Data



Immunohistochemistry analysis of paraffin-embedded Mouse brown adipose tissue using PPAR γ Rabbit pAb (A0270SP) at a dilution of 1:1600 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Rat brown adipose tissue using PPAR γ Rabbit pAb (A0270SP) at a dilution of 1:1600 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.



Confocal imaging of 3T3-L1 differentiated cells and 3T3-L1 cells using PPAR γ Rabbit pAb (A0270SP, dilution 1:1600) followed by a further incubation with Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). Objective: 100x.