# STAT5B Rabbit pAb

Catalog No.: A0275 1 Publications



## **Basic Information**

#### **Observed MW**

95kDa

#### **Calculated MW**

90kDa

#### Category

Primary antibody

#### **Applications**

ELISA,WB,IF/ICC

#### **Cross-Reactivity**

Human, Rat

## **Background**

The protein encoded by this gene is a member of the STAT family of transcription factors. In response to cytokines and growth factors, STAT family members are phosphorylated by the receptor associated kinases, and then form homo- or heterodimers that translocate to the cell nucleus where they act as transcription activators. This protein mediates the signal transduction triggered by various cell ligands, such as IL2, IL4, CSF1, and different growth hormones. It has been shown to be involved in diverse biological processes, such as TCR signaling, apoptosis, adult mammary gland development, and sexual dimorphism of liver gene expression. This gene was found to fuse to retinoic acid receptor-alpha (RARA) gene in a small subset of acute promyelocytic leukemias (APLL). The dysregulation of the signaling pathways mediated by this protein may be the cause of the APLL.

## **Recommended Dilutions**

**WB** 1:500 - 1:2000

**IF/ICC** 1:20 - 1:50

## **Immunogen Information**

**Gene ID**6777

Swiss Prot

P51692

#### Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 40-280 of human STAT5B (NP\_036580.2).

## **Synonyms**

STAT5; GHISID2; STAT5B

## **Contact**

<u>a</u>		400-999-6126
$\bowtie$		cn.market@abclonal.com.cn
•	T	www.abclonal.com.cn


## **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

#### Storage


Store at -20  $^{\circ}\text{C}.$  Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.



Western blot analysis of extracts of various cell lines, using STAT5B antibody (A0275) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.



Immunofluorescence analysis of HeLa cells using STAT5B antibody (A0275). Blue: DAPI for nuclear staining.