# MTHFR Rabbit pAb

Catalog No.: A0305



## **Basic Information**

#### **Observed MW**

75kDa

### **Calculated MW**

75kDa

### Category

Primary antibody

### **Applications**

WB,ELISA

### **Cross-Reactivity**

Rat

## **Background**

The protein encoded by this gene catalyzes the conversion of 5,10-methylenetetrahydrofolate to 5-methyltetrahydrofolate, a co-substrate for homocysteine remethylation to methionine. Genetic variation in this gene influences susceptibility to occlusive vascular disease, neural tube defects, colon cancer and acute leukemia, and mutations in this gene are associated with methylenetetrahydrofolate reductase deficiency.

## **Recommended Dilutions**

**WB** 1:500 - 1:2000

**ELISA** 

Recommended starting concentration is 1 µg/mL.
Please optimize the concentration based on your specific assay

requirements.

## Immunogen Information

**Gene ID**4524

Swiss Prot
P42898

### **Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 1-230 of human MTHFR (NP\_005948.3).

### **Synonyms**

**MTHFR** 

### **Contact**

6	400-999-6126
$\bowtie$	cn.market@abclonal.com.cn
•	www.abclonal.com.cn


### **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

### **Storage**

Store at -20  $^{\circ}\text{C}.$  Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.



Western blot analysis of lysates from Rat kidney, using MTHFR Rabbit pAb (A0305) at 1:2000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 180s.