# MEK7 Rabbit pAb

Catalog No.: A0528



### **Basic Information**

#### **Observed MW**

48kDa

#### **Calculated MW**

47kDa

#### Category

Primary antibody

#### **Applications**

WB

#### **Cross-Reactivity**

Human

### **Background**

The protein encoded by this gene is a dual specificity protein kinase that belongs to the MAP kinase kinase family. This kinase specifically activates MAPK8/JNK1 and MAPK9/JNK2, and this kinase itself is phosphorylated and activated by MAP kinase kinase kinases including MAP3K1/MEKK1, MAP3K2/MEKK2,MAP3K3/MEKK5, and MAP4K2/GCK. This kinase is involved in the signal transduction mediating the cell responses to proinflammatory cytokines, and environmental stresses. Alternative splicing results in multiple transcript variants.

## **Recommended Dilutions**

**WB** 

1:500 - 1:1000

### **Immunogen Information**

Gene ID 5609 Swiss Prot 014733

Immunogen

A synthetic peptide of human MEK7

**Synonyms** 

MEK; MKK7; JNKK2; MEK 7; MAPKK7; PRKMK7; SAPKK4; SAPKK-4; MEK7

### **Contact**

8		400-999-6126
$\bowtie$		cn.market@abclonal.com.cn
$\overline{\Box}$	ī	www.abclonal.com.cn


### **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

#### **Storage**

Store at 4°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,pH7.3.

### **Validation Data**



Western blot analysis of various lysates using MEK7 Rabbit pAb (A0528) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit  $\lg G$  (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 60s.