

SNRPA1 Rabbit mAb

Catalog No.: A0647 **Recombinant** **3 Publications**

Basic Information

Observed MW

28 kDa

Calculated MW

28 kDa

Category

Primary antibody

Applications

WB,IP,IHC-P,ELISA

Cross-Reactivity

Human, Mouse, Rat

CloneNo number

ARC2532

Background

Enables RNA binding activity. Involved in mRNA splicing, via spliceosome and spermatogenesis. Located in nuclear speck. Part of U2-type catalytic step 2 spliceosome and U2-type precatalytic spliceosome. Implicated in connective tissue disease.

Recommended Dilutions

WB 1:1000 - 1:6000

IP 0.5µg-4µg antibody for
200µg-400µg extracts of
whole cells

IHC-P 1:200 - 1:800

ELISA Recommended starting
concentration is 1 µg/mL.
Please optimize the
concentration based on
your specific assay
requirements.

Immunogen Information

Gene ID

6627

Swiss Prot

P09661

Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.


Synonyms

Lea1; U2A'; SNRPA1

Contact

 | 400-999-6126

 | cn.market@abclonal.com.cn

 | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol and 0.05% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.


Validation Data




Immunoprecipitation of SNRPA1 from 300 μ g extracts of Hep G2 cells was performed using 3 μ g of SNRPA1 Rabbit mAb (A0647). Rabbit Control IgG (AC005) was used to precipitate the Control IgG sample. IP samples were eluted with 1 \times Laemmli Buffer. The Input lane represents 10% of the total input. Western blot analysis of immunoprecipitates was conducted using SNRPA1 Rabbit mAb (A0647) at a dilution of 1:5000.




Western blot analysis of various lysates, using SNRPA1 Rabbit mAb (A0647) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25 μ g per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 3s.



Immunohistochemistry analysis of paraffin-embedded Human esophagus tissue using SNRPA1 Rabbit mAb (A0647) at a dilution of 1:200 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Human liver tissue using SNRPA1 Rabbit mAb (A0647) at a dilution of 1:200 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Human thyroid tissue using SNRPA1 Rabbit mAb (A0647) at a dilution of 1:200 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.

Validation Data



Immunohistochemistry analysis of paraffin-embedded Mouse intestine tissue using SNRPA1 Rabbit mAb (A0647) at a dilution of 1:200 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Rat intestine tissue using SNRPA1 Rabbit mAb (A0647) at a dilution of 1:200 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.