

# IGF1 Rabbit mAb, BSA and glycerol free

Catalog No.: A0830NB **Recombinant**

## Basic Information

### Observed MW

Refer to figures

### Calculated MW

22kDa

### Category

Primary antibody

### Applications

IHC-P, ELISA

### Cross-Reactivity

Human, Mouse

### CloneNo number

ARC0505

## Background

The protein encoded by this gene is similar to insulin in function and structure and is a member of a family of proteins involved in mediating growth and development. The encoded protein is processed from a precursor, bound by a specific receptor, and secreted. Defects in this gene are a cause of insulin-like growth factor I deficiency. Alternative splicing results in multiple transcript variants encoding different isoforms that may undergo similar processing to generate mature protein.

## Recommended Dilutions

**IHC-P** 1:50 - 1:200

**ELISA** Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

## Immunogen Information

### Gene ID

3479

### Swiss Prot

P05019

### Immunogen

A synthetic peptide corresponding to a sequence within amino acids 70-150 of human IGF1 (P05019).

### Synonyms

IGF; MGF; IGFI; IGF-I; IGF1

## Contact

☎ | 400-999-6126

✉ | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn)

🌐 | [www.abclonal.com.cn](http://www.abclonal.com.cn)

## Product Information

### Source

Rabbit

### Isotype

IgG

### Purification


Affinity purification

### Storage


Store at -20°C Avoid freeze / thaw cycles.  
Buffer: PBS with 0.05% proclin300, pH7.3.

## Validation Data


---



Immunohistochemistry analysis of paraffin-embedded Human liver using IGF1 Rabbit mAb, BSA and glycerol free (A0830NB) at dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate Bufferr (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Human placenta using IGF1 Rabbit mAb, BSA and glycerol free (A0830NB) at dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate Bufferr (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Mouse lung using IGF1 Rabbit mAb, BSA and glycerol free (A0830NB) at dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate Bufferr (pH 6.0) prior to IHC staining.