PPP3R1 Rabbit pAb

Catalog No.: A0954



Basic Information

Observed MW

15kDa

Calculated MW

19kDa

Category

Primary antibody

Applications

WB,IF/ICC,ELISA

Cross-Reactivity

Human, Mouse, Rat

Background

Enables cyclosporin A binding activity; phosphatase binding activity; and protein domain specific binding activity. Involved in calcineurin-NFAT signaling cascade and positive regulation of transcription by RNA polymerase II. Part of calcineurin complex. Implicated in Alzheimer's disease and dilated cardiomyopathy.

Recommended Dilutions

WB 1:500 - 1:2000

IF/ICC 1:50 - 1:200

ELISA Recommended starting concentration is 1 μg/mL.

Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID5534

Swiss Prot
P63098

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

CNB; CNB1; CALNB1; PPP3R1

Contact

a	400-999-6126
×	cn.market@abclonal.com.cr
$\overline{\Box}$	www.ahclonal.com.cr


Product Information

SourceIsotypePurificationRabbitIgGAffinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.




Western blot analysis of various lysates using PPP3R1 Rabbit pAb (A0954) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit lgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: $25\mu g$ per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 1s.



Immunofluorescence analysis of NIH/3T3 cells using PPP3R1 Rabbit pAb (A0954) at dilution of 1:100. Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.