# ERAB/HSD17B10 Rabbit mAb

Catalog No.: A0959 Recombinant 1 Publications



### **Basic Information**

### **Observed MW**

26kDa

#### **Calculated MW**

27kDa

### Category

Primary antibody

### **Applications**

WB,ELISA

### **Cross-Reactivity**

Human, Mouse, Rat

#### CloneNo number

ARC1852

# **Background**

This gene encodes 3-hydroxyacyl-CoA dehydrogenase type II, a member of the short-chain dehydrogenase/reductase superfamily. The gene product is a mitochondrial protein that catalyzes the oxidation of a wide variety of fatty acids and steroids, and is a subunit of mitochondrial ribonuclease P, which is involved in tRNA maturation. The protein has been implicated in the development of Alzheimer disease, and mutations in the gene are the cause of 17beta-hydroxysteroid dehydrogenase type 10 (HSD10) deficiency. Several alternatively spliced transcript variants have been identified, but the full-length nature of only two transcript variants has been determined.

# **Recommended Dilutions**

**WB** 1:500 - 1:1000

**ELISA** 

Recommended starting concentration is 1 µg/mL.
Please optimize the concentration based on your specific assay requirements.

## Immunogen Information

**Gene ID**3028

Swiss Prot
Q99714

### **Immunogen**

Synthetic peptide. This information is considered to be commercially sensitive.

# **Synonyms**

ABAD; CAMR; ERAB; HCD2; MHBD; HADH2; MRPP2; MRX17; MRX31; SCHAD; MRXS10; SDR5C1; HSD10MD; 17b-HSD10; DUPXp11.22; ERAB/HSD17B10

## **Contact**

6	400-999-6126
$\bowtie$	cn.market@abclonal.com.cn
•	www.abclonal.com.cn


### **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

#### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide, 0.05% BSA, 50% glycerol, pH7.3.



Western blot analysis of various lysates using ERAB/HSD17B10 Rabbit mAb (A0959) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins:  $25\mu g$  per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 10s.