

β-arrestin1 Rabbit pAb

Catalog No.: A0998

5 Publications

Basic Information

Observed MW

51kDa

Calculated MW

47kDa

Category

Primary antibody

Applications

WB, IHC-P, IF/ICC, IP, ELISA

Cross-Reactivity

Human, Mouse, Rat

Background

Members of arrestin/beta-arrestin protein family are thought to participate in agonist-mediated desensitization of G-protein-coupled receptors and cause specific dampening of cellular responses to stimuli such as hormones, neurotransmitters, or sensory signals. Arrestin beta 1 is a cytosolic protein and acts as a cofactor in the beta-adrenergic receptor kinase (BARK) mediated desensitization of beta-adrenergic receptors. Besides the central nervous system, it is expressed at high levels in peripheral blood leukocytes, and thus the BARK/beta-arrestin system is believed to play a major role in regulating receptor-mediated immune functions. Alternatively spliced transcripts encoding different isoforms of arrestin beta 1 have been described.

Recommended Dilutions

WB 1:500 - 1:1000

IHC-P 1:50 - 1:200

IF/ICC 1:50 - 1:200

IP 0.5μg-4μg antibody for
200μg-400μg extracts of
whole cellsELISA Recommended starting
concentration is 1 μg/mL.
Please optimize the
concentration based on
your specific assay
requirements.

Immunogen Information

Gene ID

408

Swiss Prot

P49407

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

ARB1; ARR1; β-arrestin1

Product Information

Source

Rabbit

Isotype

IgG

Purification

Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300, 50% glycerol, pH7.3.


Contact

 | 400-999-6126 | cn.market@abclonal.com.cn | www.abclonal.com.cn


Validation Data




Immunoprecipitation analysis of 200 µg extracts of SH-SY5Y cells using 3 µg β-arrestin1 antibody (A0998). Western blot was performed from the immunoprecipitate using β-arrestin1 (A0998) at a dilution of 1:1000.




Western blot analysis of various lysates using β-arrestin1 Rabbit pAb (A0998) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 30s.



Immunohistochemistry analysis of paraffin-embedded Rat heart using β-arrestin1 Rabbit pAb (A0998) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Mouse spleen using β-arrestin1 Rabbit pAb (A0998) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.




Immunofluorescence analysis of A-549 cells using β-arrestin1 Rabbit pAb (A0998) at dilution of 1:100 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.


Validation Data



Immunofluorescence analysis of HepG2 cells using β -arrestin1 Rabbit pAb (A0998) at dilution of 1:100 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of NIH/3T3 cells using β -arrestin1 Rabbit pAb (A0998) at dilution of 1:100 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of PC-12 cells using β -arrestin1 Rabbit pAb (A0998) at dilution of 1:100 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.