Leader in Biomolecular Solutions for Life Science



Catalog No.: A1000 4 Publications



## **Basic Information**

**Observed MW** 78kDa

**Calculated MW** 78kDa

Category Primary antibody

Applications WB, IF/ICC, IP

**Cross-Reactivity** Human, Mouse, Rat

## Background

The protein encoded by this gene is an isozyme of the long-chain fatty-acid-coenzyme A ligase family. Although differing in substrate specificity, subcellular localization, and tissue distribution, all isozymes of this family convert free long-chain fatty acids into fatty acyl-CoA esters, and thereby play a key role in lipid biosynthesis and fatty acid degradation. Several transcript variants encoding different isoforms have been found for this gene.

## **Recommended Dilutions**

## **Immunogen Information**

WB	1:200 - 1:1000	Gene ID	Swiss Prot
IF/ICC	1:50 - 1:200	2180	P33121
IP	1:50 - 1:100	Immunogen A synthetic peptide of human ACSL1	

#### **Synonyms**

ACS1; LACS; FACL1; FACL2; LACS1; LACS2; ACSL1

6

 $\sim$ 

#### 400-999-6126 Source Rabbit cn.market@abclonal.com.cn

### Ð www.abclonal.com.cn


# **Product Information**

Isotype lgG

**Purification** Affinity purification

#### Storage

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.



Western blot analysis of lysates from HepG2 cells, using ACSL1 Rabbit pAb (A1000). Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.