

TIAM1 Rabbit pAb

Catalog No.: A10252 **2 Publications**

Basic Information

Observed MW

250kDa

Calculated MW

178kDa

Category

Primary antibody

Applications

WB,IHC-P,ELISA

Cross-Reactivity

Human, Rat

Background

This gene encodes a RAC1-specific guanine nucleotide exchange factor (GEF). GEFs mediate the exchange of guanosine diphosphate (GDP) for guanosine triphosphate (GTP). The binding of GTP induces a conformational change in RAC1 that allows downstream effectors to bind and transduce a signal. This gene thus regulates RAC1 signaling pathways that affect cell shape, migration, adhesion, growth, survival, and polarity, as well as influencing actin cytoskeletal formation, endocytosis, and membrane trafficking. This gene thus plays an important role in cell invasion, metastasis, and carcinogenesis. In addition to RAC1, the encoded protein activates additional Rho-like GTPases such as CDC42, RAC2, RAC3 and RHOA. This gene encodes multiple protein isoforms that experience a diverse array of intramolecular, protein-protein, and phosphorylation interactions as well as phosphoinositide binding. Both the longer and shorter isoforms have C-terminal Dbl homology (DH) and pleckstrin homology (PH) domains while only the longer isoforms of this gene have the N-terminal myristoylation site and the downstream N-terminal PH domain, ras-binding domain (RBD), and PSD-95/DlgA/ZO-1 (PDZ) domain.

Recommended Dilutions

WB 1:500 - 1:1000

IHC-P 1:50 - 1:200

ELISA Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID

7074

Swiss Prot

Q13009

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

NEDLDS; TIAM-1; TIAM1

Contact

☎ | 400-999-6126

✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.


Validation Data



Western blot analysis of lysates from Rat testis, using TIAM1 Rabbit pAb (A10252) at 1:1000 dilution.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Enhanced Kit (RM00021).
Exposure time: 30s.



Immunohistochemistry analysis of paraffin-embedded Human colon carcinoma using TIAM1 Rabbit pAb (A10252) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Human lung cancer using TIAM1 Rabbit pAb (A10252) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.