# MZF1 Rabbit pAb

Catalog No.: A10356 2 Publications



#### **Basic Information**

#### **Observed MW**

110kDa

#### **Calculated MW**

82kDa

#### Category

Primary antibody

### **Applications**

WB,ELISA

#### **Cross-Reactivity**

Human, Mouse, Rat

## **Background**

Enables DNA-binding transcription factor activity, RNA polymerase II-specific; RNA polymerase II cis-regulatory region sequence-specific DNA binding activity; and protein homodimerization activity. Involved in negative regulation of transcription by RNA polymerase II and positive regulation of transcription by RNA polymerase II. Located in nucleoplasm.

## **Recommended Dilutions**

**WB** 1:500 - 1:2000

**ELISA** 

Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

## **Immunogen Information**

**Swiss Prot Gene ID** 7593 P28698

#### **Immunogen**

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

#### **Synonyms**

MZF-1; MZF1B; ZFP98; ZNF42; ZSCAN6; MZF1

## **Contact**

<b>a</b>		400-999-6126
$\bowtie$		cn.market@abclonal.com.cn
$\odot$	1	www.abclonal.com.cn

## **Product Information**


**Purification Source** Isotype Rabbit IgG Affinity purification

#### **Storage**

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.

## **Validation Data**



Western blot analysis of various lysates using MZF1 Rabbit pAb (A10356) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit lgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins:  $25\mu g$  per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 3s.