# **AADAC Rabbit pAb**

Catalog No.: A10365 1 Publications



### **Basic Information**

#### **Observed MW**

46kDa

#### **Calculated MW**

46kDa

#### Category

Primary antibody

#### **Applications**

ELISA,WB

#### **Cross-Reactivity**

Human, Mouse

### **Background**

Microsomal arylacetamide deacetylase competes against the activity of cytosolic arylamine N-acetyltransferase, which catalyzes one of the initial biotransformation pathways for arylamine and heterocyclic amine carcinogens

### **Recommended Dilutions**

**WB** 

1:500 - 1:2000

### **Immunogen Information**

Gene ID

Swiss Prot P22760

## 13

**Immunogen**Recombinant fusion protein containing a sequence corresponding to amino acids 170-399 of human AADAC (NP\_001077.2).

### **Synonyms**

DAC; CES5A1; AADAC

### **Contact**

6		400-999-6126
$\bowtie$		cn.market@abclonal.com.cn
•	T	www.abclonal.com.cn


### **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

#### **Storage**

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.



Western blot analysis of extracts of various cell lines, using AADAC antibody (A10365) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 30s.