# SOX4 Rabbit pAb

Catalog No.: A10717 4 Publications



## **Basic Information**

#### **Observed MW**

46kDa

#### **Calculated MW**

47kDa

### Category

Primary antibody

## **Applications**

WB,ELISA

#### **Cross-Reactivity**

Mouse

# **Background**

This intronless gene encodes a member of the SOX (SRY-related HMG-box) family of transcription factors involved in the regulation of embryonic development and in the determination of the cell fate. The encoded protein may act as a transcriptional regulator after forming a protein complex with other proteins, such as syndecan binding protein (syntenin). The protein may function in the apoptosis pathway leading to cell death as well as to tumorigenesis and may mediate downstream effects of parathyroid hormone (PTH) and PTH-related protein (PTHrP) in bone development. The solution structure has been resolved for the HMG-box of a similar mouse protein.

## **Recommended Dilutions**

**WB** 1:1000 - 1:5000

**ELISA** 

Recommended starting concentration is 1 µg/mL.
Please optimize the concentration based on your specific assay requirements.

# Immunogen Information

**Gene ID Swiss Prot** 6659 Q06945

#### **Immunogen**

Synthetic peptide. This information is considered to be commercially sensitive.

## **Synonyms**

CSS10; EVI16; SOX4

## **Contact**

| <u>a</u>                  | 400-999-6126              |
|---------------------------|---------------------------|
| ×                         | cn.market@abclonal.com.cn |
| $\overline{\mathfrak{S}}$ | www.abclonal.com.cn       |


## **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

#### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.



Western blot analysis of lysates from Mouse brain, using SOX4 Rabbit pAb (A10717) at 1:2000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 60s.