

[KO Validated] CDKN2A/p16INK4a Rabbit pAb

Catalog No.: A11058 **KO Validated**

Basic Information

Observed MW

17kDa

Calculated MW

8kDa/11kDa/12kDa/13kDa/16kDa/17kDa

Category

Primary antibody

Applications

ELISA,WB,IP

Cross-Reactivity

Human

Background

This gene generates several transcript variants which differ in their first exons. At least three alternatively spliced variants encoding distinct proteins have been reported, two of which encode structurally related isoforms known to function as inhibitors of CDK4 kinase. The remaining transcript includes an alternate first exon located 20 Kb upstream of the remainder of the gene; this transcript contains an alternate open reading frame (ARF) that specifies a protein which is structurally unrelated to the products of the other variants. This ARF product functions as a stabilizer of the tumor suppressor protein p53 as it can interact with, and sequester, the E3 ubiquitin-protein ligase MDM2, a protein responsible for the degradation of p53. In spite of the structural and functional differences, the CDK inhibitor isoforms and the ARF product encoded by this gene, through the regulatory roles of CDK4 and p53 in cell cycle G1 progression, share a common functionality in cell cycle G1 control. This gene is frequently mutated or deleted in a wide variety of tumors, and is known to be an important tumor suppressor gene.

Recommended Dilutions

WB 1:500 - 1:1000

IP 0.5µg-4µg antibody for
200µg-400µg extracts of
whole cells

Immunogen Information

Gene ID

1029

Swiss Prot

P42771/Q8N726

Immunogen

A synthetic peptide corresponding to a sequence within amino acids 1-156 of human CDKN2A/p16INK4a (NP_000068.1).

Synonyms

ARF; MLM; P14; P16; P19; CMM2; INK4; MTS1; TP16; CDK4i; CDKN2; INK4A; MTS-1; P14ARF; P19ARF; P16INK4; P16INK4A; P16-INK4A; 4a

Contact

☎ | 400-999-6126

✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.


Validation Data



Immunoprecipitation analysis of 200 µg extracts of 293T cells, using 3 µg CDKN2A/p16INK4a antibody (A11058). Western blot was performed from the immunoprecipitate using CDKN2A/p16INK4a antibody (A11058) at a dilution of 1:1000.



Western blot analysis of lysates from wild type (WT) and CDKN2A/p16INK4a knockout (KO) 293T cells, using [KO Validated] CDKN2A/p16INK4a Rabbit pAb (A11058) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 180s.



Western blot analysis of lysates from HeLa cells, using [KO Validated] CDKN2A/p16INK4a Rabbit pAb (A11058) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 180s.