Galectin 3/LGALS3 Rabbit mAb

Catalog No.: A11198 Recombinant 5 Publications



Basic Information

Observed MW

28kDa

Calculated MW

26kDa

Category

Primary antibody

Applications

WB,IHC-P,IP,ELISA

Cross-Reactivity

Human, Mouse, Rat

CloneNo number

ARC0542

Background

This gene encodes a member of the galectin family of carbohydrate binding proteins. Members of this protein family have an affinity for beta-galactosides. The encoded protein is characterized by an N-terminal proline-rich tandem repeat domain and a single C-terminal carbohydrate recognition domain. This protein can self-associate through the N-terminal domain allowing it to bind to multivalent saccharide ligands. This protein localizes to the extracellular matrix,the cytoplasm and the nucleus. This protein plays a role in numerous cellular functions including apoptosis,innate immunity,cell adhesion and T-cell regulation. The protein exhibits antimicrobial activity against bacteria and fungi. Alternate splicing results in multiple transcript variants.

Recommended Dilutions

WB 1:1000 - 1:6000

IHC-P 1:200 - 1:800

IP 0.5μg-4μg antibody for

200μg-400μg extracts of

whole cells

ELISA Recommended starting

concentration is 1 µg/mL.

Please optimize the
concentration based on
your specific assay
requirements.

Immunogen Information

Gene ID3958

Swiss Prot
P17931

Immunogen

A synthetic peptide corresponding to a sequence within amino acids 1-100 of human Galectin 3/LGALS3 (P17931).

Synonyms

L31; GAL3; MAC2; CBP35; GALBP; GALIG; LGALS2; Galectin 3/LGALS3

Contact

6		400-999-6126
\bowtie		cn.market@abclonal.com.cn
\odot	ī	www.abclonal.com.cn


Product Information

SourceIsotypePurificationRabbitIgGAffinity purification


Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide, 0.05% BSA, 50% glycerol, pH7.3.



Immunoprecipitation analysis of 300 μ g extracts of HT-29 cells using 3 μ g Galectin 3/LGALS3 antibody (A11198). Western blot was performed from the immunoprecipitate using Galectin 3/LGALS3 antibody (A11198) at a dilution of 1:1000.




Western blot analysis of various lysates using Galectin 3/LGALS3 Rabbit mAb (A11198) at 1[]1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.


Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).


Exposure time: 10s.



Immunohistochemistry analysis of paraffinembedded Human colon tissue using Galectin 3/LGALS3 Rabbit mAb (A11198) at a dilution of 1:500 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.




Immunohistochemistry analysis of paraffinembedded Human pancreas tissue using Galectin 3/LGALS3 Rabbit mAb (A11198) at a dilution of 1:500 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.




Immunohistochemistry analysis of paraffinembedded Human tonsil tissue using Galectin 3/LGALS3 Rabbit mAb (A11198) at a dilution of 1:500 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.



Immunohistochemistry analysis of paraffinembedded Rat colon tissue using Galectin 3/LGALS3 Rabbit mAb (A11198) at a dilution of 1:500 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.



Immunohistochemistry analysis of paraffinembedded Rat liver tissue using Galectin 3/LGALS3 Rabbit mAb (A11198) at a dilution of 1:500 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.



Immunohistochemistry analysis of paraffinembedded Rat spleen tissue using Galectin 3/LGALS3 Rabbit mAb (A11198) at a dilution of 1:500 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.