# PIK3R3/p55PIK Rabbit mAb

Catalog No.: A11260 Recombinant 1 Publications



## **Basic Information**

### **Observed MW**

54kDa

### **Calculated MW**

54kDa

### Category

Primary antibody

### **Applications**

ELISA,WB,IHC-P,IF/ICC

#### **Cross-Reactivity**

Human, Mouse, Rat

## **Background**

Phosphatidylinositol 3-kinase (PI3K) phosphorylates phosphatidylinositol and similar compounds, which then serve as second messengers in growth signaling pathways. PI3K is composed of a catalytic and a regulatory subunit. The protein encoded by this gene represents a regulatory subunit of PI3K. The encoded protein contains two SH2 domains through which it binds activated protein tyrosine kinases to regulate their activity.

# **Recommended Dilutions**

WB	1:500 - 1:1000
IHC-P	1:50 - 1:200
IF/ICC	1:50 - 1:200

# **Immunogen Information**

**Gene ID Swiss Prot**8503
Q92569

#### **Immunogen**

Recombinant protein of human PIK3R3

### **Synonyms**

p55; p55PIK; p55-GAMMA; PIK3R3/p55PIK

### **Contact**

6		400-999-6126
$\bowtie$		cn.market@abclonal.com.cn
•	T	www.abclonal.com.cn


### **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide, 0.05% BSA, 50% glycerol, pH7.3.



Western blot analysis of various lysates using PIK3R3/p55PIK Rabbit mAb (A11260). Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins:  $25\mu g$  per lane.

Blocking buffer: 3% nonfat dry milk in TBST.