

TRADD Rabbit pAb

Catalog No.: A1145 **2 Publications**

Basic Information

Observed MW

36kDa

Calculated MW

34kDa

Category

Primary antibody

Applications

WB, ELISA

Cross-Reactivity

Human, Mouse

Background

Predicted to enable death domain binding activity; identical protein binding activity; and signaling receptor binding activity. Involved in positive regulation of hair follicle development. Acts upstream of or within extrinsic apoptotic signaling pathway. Located in cytoplasm. Is expressed in adrenal gland; genitourinary system; liver; lung; and spleen. Orthologous to human TRADD (TNFRSF1A associated via death domain).

Recommended Dilutions

WB 1:1000 - 1:5000

ELISA Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID

8717

Swiss Prot

Q15628

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 199-312 of human TRADD (NP_003780.1).

Synonyms

9130005N23Rik; TRADD

Contact

☎ | 400-999-6126

✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH 7.3.

Validation Data



Western blot analysis of lysates from MCF7 cells, using TRADD Rabbit pAb (A1145) at 1:2000 dilution.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 90s.