

# TRADD Rabbit pAb

Catalog No.: A1145

1 Publications

## Basic Information

### Observed MW

36kDa

### Calculated MW

34kDa

### Category

Primary antibody

### Applications

ELISA, WB

### Cross-Reactivity

Human, Mouse

## Background

Predicted to enable death domain binding activity; identical protein binding activity; and signaling receptor binding activity. Involved in positive regulation of hair follicle development. Acts upstream of or within extrinsic apoptotic signaling pathway. Located in cytoplasm. Is expressed in adrenal gland; genitourinary system; liver; lung; and spleen. Orthologous to human TRADD (TNFRSF1A associated via death domain).

## Recommended Dilutions

WB 1:1000 - 1:5000

## Immunogen Information

### Gene ID

8717

### Swiss Prot

Q15628

### Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 199-312 of human TRADD (NP\_003780.1).

### Synonyms

9130005N23Rik; TRADD

## Contact

☎ | 400-999-6126

✉ | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn)

🌐 | [www.abclonal.com.cn](http://www.abclonal.com.cn)

## Product Information

### Source

Rabbit

### Isotype

IgG

### Purification

Affinity purification


### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH 7.3.

## Validation Data

---



Western blot analysis of lysates from MCF7 cells, using TRADD Rabbit pAb (A1145) at 1:2000 dilution.  
Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.  
Lysates/proteins: 25µg per lane.  
Blocking buffer: 3% nonfat dry milk in TBST.  
Detection: ECL Basic Kit (RM00020).  
Exposure time: 90s.