# MC3R Rabbit pAb

Catalog No.: A11609



# **Basic Information**

#### **Observed MW**

36kDa

### **Calculated MW**

36kDa

### Category

Primary antibody

## **Applications**

ELISA,WB

#### **Cross-Reactivity**

Mouse, Rat

# **Background**

This gene encodes a G-protein-coupled receptor for melanocyte-stimulating hormone and adrenocorticotropic hormone that is expressed in tissues other than the adrenal cortex and melanocytes. This gene maps to the same region as the locus for benign neonatal epilepsy. Mice deficient for this gene have increased fat mass despite decreased food intake, suggesting a role for this gene product in the regulation of energy homeostasis. Mutations in this gene are associated with a susceptibility to obesity in humans.

## **Recommended Dilutions**

**WB** 

1:500 - 1:2000

# **Immunogen Information**

Gene ID 4159 **Swiss Prot** 

P41968

#### **Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 1-120 of human MC3R (NP\_063941.3).

## **Synonyms**

MC3; OB20; OQTL; BMIQ9; MC3-R; MC3R

## **Contact**

| <b>a</b>  |   | 400-999-6126              |
|-----------|---|---------------------------|
| $\bowtie$ |   | cn.market@abclonal.com.cn |
| •         | Ī | www.abclonal.com.cn       |


## **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

## **Storage**

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.



Western blot analysis of various lysates using MC3R Rabbit pAb (A11609) at 1:3000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 30s.