

# OCIAD1 Rabbit pAb

**Catalog No.: A11630**

## Basic Information

### Observed MW

28kDa

### Calculated MW

28kDa

### Category

Primary antibody

### Applications

ELISA, WB, IF/ICC

### Cross-Reactivity

Human, Mouse

## Background

Predicted to be involved in several processes, including hematopoietic stem cell homeostasis; positive regulation of receptor signaling pathway via JAK-STAT; and regulation of stem cell differentiation. Located in membrane.

## Recommended Dilutions

**WB** 1:500 - 1:2000

**IF/ICC** 1:50 - 1:200

## Immunogen Information

### Gene ID

54940

### Swiss Prot

Q9NX40

### Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 1-245 of human OCIAD1 (NP\_060300.1).

### Synonyms

OCIA; ASRIJ; TPA018; OCIAD1

## Contact

 | 400-999-6126

 | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn)

 | [www.abclonal.com.cn](http://www.abclonal.com.cn)

## Product Information

### Source

Rabbit

### Isotype

IgG

### Purification

Affinity purification


### Storage

Store at -20°C. Avoid freeze / thaw cycles.


Buffer: PBS with 0.01% thimerosal, 50% glycerol, pH7.3.

## Validation Data

---



Western blot analysis of lysates from mouse liver, using OCIAD1 Rabbit pAb (A11630) at 1:3000 dilution.  
Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.  
Lysates/proteins: 25µg per lane.  
Blocking buffer: 3% nonfat dry milk in TBST.  
Detection: ECL Basic Kit (RM00020).  
Exposure time: 30s.



Confocal immunofluorescence analysis of U2OS cells using OCIAD1 Rabbit pAb (A11630) at dilution of 1:100. Blue: DAPI for nuclear staining.