

CHAMP1 Rabbit pAb

Catalog No.: A11690

Basic Information

Observed MW

130kDa

Calculated MW

89kDa

Category

Primary antibody

Applications

ELISA, WB, IF/ICC

Cross-Reactivity

Human, Mouse

Background

This gene encodes a zinc finger protein that functions as a regulator of chromosome segregation in mitosis. The encoded protein is required for correct alignment of chromosomes on the metaphase plate, and plays a role in maintaining the attachment of sister kinetochores to microtubules from opposite spindle poles. Mutations in this gene are associated with an autosomal dominant form of intellectual disability.

Recommended Dilutions

WB	1:500 - 1:2000
IF/ICC	1:50 - 1:100

Immunogen Information

Gene ID

283489

Swiss Prot

Q96JM3

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 563-812 of human CHAMP1 (NP_115812.1).

Synonyms

CAMP; CHAMP; MRD40; ZNF828; C13orf8; NEDHILD; CHAMP1

Contact

☎ | 400-999-6126

✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal, 50% glycerol, pH7.3.

Validation Data



Western blot analysis of various lysates using CHAMP1 Rabbit pAb (A11690) at 1:3000 dilution.
Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 1s.



Immunofluorescence analysis of HeLa cells using CHAMP1 Rabbit pAb (A11690) at dilution of 1:100 (40x lens). Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.