TDP-43/TARDBP Rabbit pAb

Catalog No.: A1183 1 Publications



Basic Information

Observed MW

42kDa

Calculated MW

45kDa

Category

Primary antibody

Applications

WB,IHC-P,IP,ELISA,RIP

Cross-Reactivity

Human, Mouse, Rat

Background

HIV-1, the causative agent of acquired immunodeficiency syndrome (AIDS), contains an RNA genome that produces a chromosomally integrated DNA during the replicative cycle. Activation of HIV-1 gene expression by the transactivator Tat is dependent on an RNA regulatory element (TAR) located downstream of the transcription initiation site. The protein encoded by this gene is a transcriptional repressor that binds to chromosomally integrated TAR DNA and represses HIV-1 transcription. In addition, this protein regulates alternate splicing of the CFTR gene. A similar pseudogene is present on chromosome 20.

Recommended Dilutions

WB 1:500 - 1:2000

IHC-P 1:50 - 1:200

IP 0.5μg-4μg antibody for

200μg-400μg extracts of

whole cells

ELISA Recommended starting

concentration is 1 µg/mL.

Please optimize the concentration based on your specific assay requirements.

RIP 1:20 - 1:50

Contact

<u>a</u>	400-999-6126
\bowtie	cn.market@abclonal.com.cn
•	www.abclonal.com.cn

Immunogen Information

Gene ID23435

Swiss Prot
Q13148

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

ALS10; TDP-43; TDP-43/TARDBP


Product Information

SourceIsotypePurificationRabbitIgGAffinity purification


Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.




Immunoprecipitation analysis of 200 μg extracts of SW620 cells using 3 μg TDP-43/TARDB antibody (A1183). Western blot was performed from the immunoprecipitate using TDP-43/TARDB antibody (A1183) at a dilution of 1:500.




Western blot analysis of various lysates using TDP-43/TARDB Rabbit pAb (A1183) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit lgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane.


Blocking buffer: 3% nonfat dry milk in TBST.



Immunohistochemistry analysis of paraffinembedded Rat spleen using TDP-43/TARDB Rabbit pAb (A1183) at dilution of 1:200 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.



Immunohistochemistry analysis of paraffinembedded Human gastric cancer using TDP-43/TARDB Rabbit pAb (A1183) at dilution of 1:200 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.



Immunohistochemistry analysis of paraffinembedded Mouse brain using TDP-43/TARDB Rabbit pAb (A1183) at dilution of 1:200 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.