β-Tubulin Rabbit mAb

Catalog No.: A12289 Recombinant 52 Publications



Basic Information

Observed MW

55 kDa

Calculated MW

50 kDa

Category

Primary antibody

Applications

WB,IHC-P,IP,ELISA

Cross-Reactivity

Human, Mouse, Rat

CloneNo number

ARC0203

Background

This gene encodes a beta tubulin protein. This protein forms a dimer with alpha tubulin and acts as a structural component of microtubules. Mutations in this gene cause cortical dysplasia, complex, with other brain malformations 6. Alternative splicing results in multiple splice variants. There are multiple pseudogenes for this gene on chromosomes 1, 6, 7, 8, 9, and 13.

Recommended Dilutions

WB 1:3000-1:30000

IHC-P 1:50 - 1:200

IP 0.5μg-4μg antibody for 200μg-400μg extracts of

whole cells

ELISA Recommended starting

concentration is 1 µg/mL.

Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID203068

Swiss Prot
P07437

Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms

M40; TUBB1; TUBB5; CDCBM6; CSCSC1; OK/SW-cl.56; β -Tubulin

Contact

6		400-999-6126
\bowtie		cn.market@abclonal.com.cn
\odot	ī	www.abclonal.com.cn


Product Information

SourceIsotypePurificationRabbitIgGAffinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.09% Sodium azide, 0.05% BSA, 50% glycerol, pH7.3.




Western blot analysis of various lysates using β -Tubulin Rabbit mAb (A12289) at 1:5000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25 μ g per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 20s.



Western blot analysis of lysates from 293F cells, using β -Tubulin Rabbit mAb (A12289) at 1:1000-1:30000 dilution.


Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.


Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 10s.



Immunoprecipitation analysis of 300 μg extracts of 293T cells using 3 μg β -Tubulin Rabbit mAb (A12289). Western blot was performed from the immunoprecipitate using β -Tubulin Rabbit mAb (A12289) at a dilution of 1:1000.



Validation Data

Immunohistochemistry analysis of paraffinembedded Rat kidney tissue using $\beta\text{-}\text{Tubulin}$ Rabbit mAb (A12289) at a dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.